



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Crystal structure of 4-methoxycarbonyl-1,3-dithiole-2-thione, C₅H₄O₂S₃

Citation for published version:

Darviche, F, Asghari, G-R, Parsapour, M, Balalaie, S, Rominger, F & Robertson, N 2007, 'Crystal structure of 4-methoxycarbonyl-1,3-dithiole-2-thione, C₅H₄O₂S₃', *Zeitschrift für Kristallographie - New Crystal Structures*, vol. 222, no. 1, pp. 27-28. <https://doi.org/10.1524/ncrs.2007.0010>

Digital Object Identifier (DOI):

[10.1524/ncrs.2007.0010](https://doi.org/10.1524/ncrs.2007.0010)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Zeitschrift für Kristallographie - New Crystal Structures

Publisher Rights Statement:

Copyright © 2014 Oldenbourg Wissenschaftsverlag GmbH. This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 License.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Crystal structure of 4-methoxycarbonyl-1,3-dithiole-2-thione, $C_5H_4O_2S_3$


Fatemeh Darviche^{*,I}, Gholam-Reza Asghari^I, Melika Parsapour^I, Saeed Balalaie^I, Frank Rominger^{II} and Neil Robertson^{III}

^I K.N. Toosi University of Technology, Chemistry Department, P.O. Box 15875-4416, Tehran, Iran

^{II} Universität Heidelberg, Organisch-Chemisches Institut, Im Neuenheimer Feld 270, 69120 Heidelberg, Germany

^{III} University of Edinburgh, School of Chemistry, King's buildings, West Mains Road, Edinburgh EH9 3JJ, UK

Received November 6, 2006, accepted and available on-line March 23, 2007; CCDC no. 1267/1927



Abstract

$C_5H_4O_2S_3$, monoclinic, $P12_1/c1$ (no. 14), $a = 3.9797(1)$ Å, $b = 6.0394(2)$ Å, $c = 31.1165(6)$ Å, $\beta = 91.868(1)^\circ$, $V = 747.5$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.119$, $wR_{\text{ref}}(F^2) = 0.324$, $T = 200$ K.

Source of material

In the literature 4-methoxycarbonyl-1,3-dithiole-2-thione was obtained in three steps [1]. In the present one-pot synthesis the title compound was obtained under solvent free conditions catalyzed by potassium cyanide in the presence of silica as solid support at 200 °C (85 % yield). 0.2 mmol of 4,5-bis(carbonyl-methoxy)-1,3-dithiole-2-thione, 0.4 mmol of potassium cyanide and 0.2 g of silica were mixed in a mortar. This powder was heated for 1 h at 200 °C. The color of the mixture was turned from yellow to light brown. The pure compound was obtained by filtration of the solution in dichloromethane, followed by evaporation of solvent and preparative chromatography over silica gel. The product was characterized by IR, ¹H NMR, and ¹³C NMR spectroscopy.

Experimental details

The X-ray reflections were rather broad and diffuse, but the data turned out to be useable. All hydrogen atoms were treated using appropriate riding models. The bad quality of the crystals and the dataset were the reasons for the large R values.

Discussion

Ever since the discovery of the first metallic charge transfer complex composed of TTF (tetrathiafulvalene) and TCNQ (tetracyanoquinodimethane) [2], modifications of the TTF skeleton have received considerable attention from synthetic chemists in exploration of new molecular-based organic metals. Phosphite cross-couplings for the preparation of functionalized TTF derivatives were first investigated because this method is generally simple when yields are good and when products are separable [3–6]. Thus the interest in the synthesis of various 1,3-dithiole-2-chalcogenone is evident, and promoted us to take up this project. As expected the interatomic distances determined in the 1,3-dithiole-2-thione unit are all similar to analogous values known from the literature. The heterocyclic five-membered dithiolethione ring is planar, and also the carboxyl group stands in the same plane, thus is completely in conjugation. The crystal packing is dominated by very short sulfur...sulfur contacts involving two of the three sulfur atoms and building two-dimensional double layers of molecules. The S1...S1 contacts with 3.51 Å, the S1...S2 contacts with 3.55 Å, and the S2...S2 contacts with 3.42 Å are much shorter than the sum of the van der Waals radii of 3.7 Å [8].

Table 1. Data collection and handling.

Crystal:	yellow polyhedron, size 0.05 × 0.24 × 0.29 mm
Wavelength:	Mo K_α radiation (0.71073 Å)
μ :	9.21 cm ⁻¹
Diffractionmeter, scan mode:	Bruker SMART CCD, ϕ/ω
$2\theta_{\text{max}}$:	51.4°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	6320, 1419
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2\sigma(I_{\text{obs}})$, 1266
$N(\text{param})_{\text{refined}}$:	92
Program:	SHELXTL [9]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U_{iso}
H(2)	4e	0.6598	0.8751	0.4062	0.026
H(5A)	4e	1.1270	0.8647	0.2795	0.054
H(5B)	4e	0.9949	1.1074	0.2908	0.054
H(5C)	4e	0.7506	0.9358	0.2661	0.054

* Correspondence author (e-mail: darvish@kntu.ac.ir)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
S(1)	4e	0.0990(8)	0.1705(5)	0.4564(1)	0.029(2)	0.012(1)	0.044(2)	−0.008(1)	0.005(1)	0.001(1)
S(2)	4e	0.3996(7)	0.6179(4)	0.45127(9)	0.031(2)	0.007(1)	0.034(2)	−0.004(1)	0.004(1)	−0.002(1)
S(3)	4e	0.3851(8)	0.3412(5)	0.3752(1)	0.033(2)	0.009(1)	0.035(2)	−0.001(1)	0.002(1)	−0.004(1)
C(1)	4e	0.282(3)	0.364(2)	0.4288(4)	0.019(5)	0.004(5)	0.041(6)	−0.002(4)	0.001(5)	0.003(4)
C(2)	4e	0.569(3)	0.730(2)	0.4060(4)	0.020(6)	0.011(5)	0.035(6)	−0.004(4)	0.002(4)	0.002(4)
C(3)	4e	0.567(3)	0.605(2)	0.3703(4)	0.019(5)	0.006(5)	0.038(6)	0.001(4)	−0.001(4)	0.001(4)
C(4)	4e	0.687(3)	0.661(2)	0.3282(4)	0.020(6)	0.026(6)	0.033(6)	0.004(5)	−0.001(5)	0.001(5)
O(4)	4e	0.675(2)	0.547(2)	0.2971(3)	0.037(5)	0.028(5)	0.039(5)	0.002(4)	0.004(4)	−0.004(4)
O(5)	4e	0.818(2)	0.871(2)	0.3284(3)	0.045(5)	0.021(5)	0.038(5)	0.001(4)	0.005(4)	0.003(4)
C(5)	4e	0.932(3)	0.951(2)	0.2880(4)	0.035(7)	0.031(7)	0.042(8)	−0.007(6)	0.002(6)	0.009(6)

References

1. Pittman, C. U. Jr.; Narita, M.; Liang, Y. F.: Preparation and Reactions of Bi-functionalized Tetrathiafulvalenes. *J. Org. Chem.* **41** (1976) 2855-2860.
2. Ferraris, J. P.; Cowan, D. O.; Walatka, V.; Perlstein, J. H.: Electron Transfer in a New Highly Conducting Donor-Acceptor Complex. *J. Am. Chem. Soc.* **95** (1973) 948-949.
3. Hartzler, H. D.: Nucleophilic 1,3-Dithiolium Carbene. *J. Am. Chem. Soc.* **92** (1970) 1412-1413.
4. Hurley, M. F.; Chambers, J. Q.: Electrochemical Reduction of Carbon Disulfide. Synthesis of Carbon Sulfide Heterocycles. *J. Org. Chem.* **46** (1981) 778-782.
5. Krief, A.: Syntheses of Tetraheterofulvalenes and of Vinylene Trihetero-carbonates Strategy and Practice. *Tetrahedron* **42** (1986) 1204-1252.
6. Kini, A. H.; Tytko, S. F.; Hunt, J. E.; Williams, J. M.: 4,5-Methylenedithio-4',5'-propylenedithiotetrathiafulvalene (MPT) and 4,5-ethylenedithio-4',5'-propylenedithiotetrathiafulvalene (EPT). *Tetrahedron Lett.* **28** (1987) 4153-4156.
7. Otsubo, T.; Aso, Y.; Takimiya, K.: Dimeric Tetrathiafulvalenes: New electron donors. *Adv. Mater.* **8** (1996) 203-211.
8. Werz, D. B.; Staeb, T. H.; Benisch, C.; Rausch, B. J.; Rominger, F.; Gleiter, R.: Self-Organisation of Chalcogen-Containing Cyclic Alkynes and Alkenes To Yield Columnar Structures. *Org. Lett.* **4** (2002) 339-342.
9. Sheldrick, G. M.: SHELXTL. Structure Determination Software Package. V6.12. Bruker AXS, Madison, Wisconsin, USA 2001.